

## ERRATUM

In UHM 2005, 32(2):85-88, the “Reference” section which appears on page 87-88 is incorrect. It should read as follows:

### REFERENCES

1. Marroni A, Bennett PB, Cronje FJ *et al.* A deep stop during decompression from 82 fsw (25 m) significantly reduces bubbles and fast tissue gas tensions. *Undersea Hyperb Med* 2004; 31:233-43.
2. Sawatzky KD, Nishi RY. Assessment of inter-rater agreement on the grading of intravascular bubble signals. *Undersea Biomed Res* 1991; 18:373-96.
3. Nishi RY, Brubakk AO, Eftedal OS. Bubble detection. In: Brubakk AO, Neuman TS, Eds. Bennett and Elliott's physiology and medicine of diving. 5th edition. London: Saunders, 2003: 501-29.
4. Moon RE, Camporesi EM, Kisslo JA. Patent foramen ovale and decompression sickness in divers. *Lancet* 1989; 1:513-4.
5. Wilmshurst P, Bryson P. Relationship between the clinical features of neurological decompression illness and its causes. *Clin Sci (Lond)* 2000; 99:65-75.
6. Marroni A. Use of a deep stop during the ascent from 25 metres could significantly reduce the incidence of DCI of the neurological type in recreational diving. *Alert Diver* 2004; 147

The Managing Editor apologizes for the error.